

Complications of appendicular peritonitis in girls

Azamat Muxitdinovich Shamsiev
Shukhrat Abdurasulovich Yusupov
Azizbek Utkirovich Kiyamov
Anvar Komilovich Rakhimov
Samarkand State Medical University

Abstract: After the operation for gangrenous - perforated appendicitis, miscarriage occurs in 11%, infertility in 62% of cases. 154 girls aged 3 to 15 years were examined. For the purpose of diagnosis, a complex of examinations was carried out: collection of anamnesis data, analysis of menstrual and generative functions, bimanual examination, colposcopy, microbiological examination of the contents of the cervical canal and vagina, polymerase chain reaction to reveal sexually transmitted diseases, assessment of the functional state of the ovaries, examination of the sexual partner, hormonal examination, ultrasonic examination of small pelvis organs and thyroid gland, examination of the mammary glands. The developed algorithm makes it possible to carry out timely diagnosis and treatment of complications of reproductive function in girls after suffering from prevalent appendicular peritonitis.

Keywords: girls, prevalent appendicular peritonitis, reproductive function

Relevance. It is known that genital and extragenital diseases undergone in childhood are a favourable factor for the development of many gynecological diseases in an adultwomen. Acute and chronic inflammations of the genitals, especially of the fallopian tubes (20%) are of essential significance among female infertility causes. In about 40-85% of cases, the cause of this is the so-called tubal-peritoneal factor developed after the acute and chronic inflammatory process of uterine appendages, prior operations in the area of the small pelvis and adjacent abdominal organs. The tubal-peritoneal factor is primarily associated with inflammatory changes, the formation of adhesions in the small pelvis cavity, accompanied by impaired patency and functional viability of the fallopian tubes. And even after the surgical removal of the adhesive process in the peritubular space and the restoration of permeability of the fallopian tubes, pregnancy occurs only in 30% of cases. It is caused by recurrence of adhesive formation making 80-90% [1,3,5,7,9] according to the data of various authors.

Acute peritonitis is one of the most serious complications caused by the diseases of the abdominal organs. Acute destructive appendicitis takes the first place among the causes of acute peritonitis. In this case, the girls' organs of small

pelvis, the uterus and its appendages are also involved in the pathological process. An increase in incidence of appendicitis, statistically coincides with the second phase of puberty, in which all sections of the reproductive system are highly susceptible to pathological effects, that complicates the problem furthermore. Inflammation of the internal genital organs also leads to the disruption of menstrual activity. In this case, the central section, which regulates menstruation, is rarely involved in the pathological process. Menstrual irregularities during puberty (irregular menstruation, hypomenstrual syndrome, secondary amenorrhea) more often occur after catarrhal appendicitis (83%), and less often in gangrenous and gangrenous-perforated appendicitis (70%). The results of the analysis of gynecological diseases of women who underwent appendectomy in childhood, revealed chronic adnexitis (17%) menstrual irregularities (39%), primary infertility (25,4%), complicated course of pregnancy (33,7%), and childbirth (30,6%) that indicated a high rate compared to this group of the population. After surgery for gangrenous - perforated appendicitis, miscarriage occurs in 11%, infertility in 62% of cases [2, 4, 6, 9, 10].

The purpose of the study: the aim of the research is to develop and put into practice an algorithm for monitoring the activity of the reproductive system in girls and young women who underwent prevalent appendicular (PAP) peritonitis in childhood.

Materials and research methods: 154 women, who had undergone widespread appendicular peritonitis at the age of 3 to 15 years and had been operated in the 2nd Clinic of Samarkand State Medical Institute, were examined. During the operation, phlegmonous appendicitis was diagnosed in 3 (1,9%) patients, gangrenous appendicitis in 19 (12,4%) and gangrenous perforated appendicitis in 74 (85,7%). Their distribution according to prevalence of peritonitis was the following: diffuse peritonitis was identified in 82 (53,2%) prevalent peritonitis in 72 (46,8%) patients. Patients were discharged in satisfactory condition and were followed up in a dispensary.

All of those patients who had undergone PAP in their childhood were performed a complex of clinical-laboratory-instrumental examination for diagnostic aim which included: Careful collection of anamnestic data, analysis of menstrual and generative function, bimanual examination, colposcopy, microbiological examination of contents from the cervical canal and vagina, polymerase-chain reaction to reveal sexually transmitted diseases, assessment of the functional condition of the ovaries, hormonal examination, ultrasonic examination (US) of small pelvis organs and thyroid gland, examination of mammary glands.

Of particular importance in the evaluation of reproductive function was the condition of the menstrual cycle in the patients examined. We have determined

the degree of sexual development of patients in both groups. According to the period of sexual development, the examined patients were divided into 2 age categories - 12-15 years, as well as 16 years and older.

The assessment of sexual development included the degree of development of the mammary glands (Ma), the hair-covering on the pubic (P) and axillary (Ax) areas, the terms of the first menstruation (Me) and was calculated by the formula: $Max + Px + Axx + Mex$. The results of the study were statistically processed using a software package Microsoft Office Excel-2012 based on built-in functions.

Results and discussion: in the acute period before the operation menarche was observed in 26 (16,9%) girls, and the operation could no longer affect the terms of appearance of the first menstruation in these patients, therefore their data were not taken into account in the analysis of the effect of undergone peritonitis on the formation of the menstrual cycle.

In patients most often (63,0%) menarche occurred at the age of 13-14 years, but in case of disturbance of reproductive function menstruation is delayed. By the age of 12, the first menstruation was noted in 11,1% of cases in the group, by the age of 15 in 17,8%.

The nature of the post-operative course of PAP has a statistically significant effect on menstruation tenderness and on the frequency of pregnancy, as well as on the nature of its course. Thus, 38,7% of patients more often complained of painful menstruation. In addition, in 48,4% of cases, ectopic pregnancies, miscarriages and premature births were observed, and in 30,5% infertility development was noted.

In 24,4% of cases with pathology resulting in problems of reproductive function in women who had undergone peritonitis surgery in childhood, cystic ovarian changes and the adhesive process prevailed.

The adhesive process was observed in 26,7%. During the examination, the adhesive process was often accompanied by retroflexion of the uterus, which is also one of the causes of infertility. Adhesions often accompany inflammatory diseases of pelvic organs. Thus, among the cases of inflammatory diseases of the genitals without signs of endometriosis, 85,3% of the patients had adhesions and only 14,7% had no adhesions.

In 23 (92,0%) women with infertility, of 25 after PAP in childhood, hysterosalpingography was performed to exclude or confirm tubal-peritoneal infertility. In 2 other patients, the absence of ovulation (1) and uterine hypoplasia (1) were diagnosed. In our observations, the frequency of development of occlusal lesions of the fallopian tubes in women after the previous PAP was 11 (47,8%) cases, while almost all patients - 10 (90,9%) - had a lesion of the right fallopian

tube, including 2 with bilateral lesion, 4 with full occlusion on the right and 4 more with partial occlusion on the right. An isolated lesion of the left fallopian tube was diagnosed in only 1 case in the form of partial occlusion. It is interesting to note that in infertile women with intact permeability of both fallopian tubes, we often found polycystic ovary disease (3), and in 1 case, uterine hypoplasia was revealed.

We analyzed the level of pituitary hormones (FSH and LH), as well as progesterone and estradiol in phases I and II of the menstrual cycle. The data in Table 1 show that in patients who had undergone PAP in childhood, hormone levels were slightly higher relative to the reference values. High or on the contrary moderately low levels of pituitary hormones compared to the norm are permissible in different phases of the menstrual cycle.

When comparing ovarian hormone levels, estradiol and progesterone were found to be within the normal limits in patients suffering from PAP.

PAP treatment, as you know, involves the use of large doses of powerful broad-spectrum antibiotics, often using their combination with antifungal agents. Their use in itself contributes to the development of intra-abdominal adhesions. Ozone therapy proposed in our work makes it possible to fight effectively against pathogenic intra-abdominal microflora, and this fact permits to significantly terms reduce the and doses of antibiotics, and to optimize their use.

The prediction of pregnancy depends on the complex effect of many factors:

- Pre-operative training in the form of adequate rehabilitation of all infection foci.
- The volume of performed intervention, preservation of fimbrial sections of uterine tubes and their epiteal covering, necessity to carry out combined reconstructive and plastic operations.
- The course of postoperative period. Step-by-step rehabilitation promotes spontaneous pregnancy in 10-20% of patients.
- Hysterosalpingography - diagnosis of the presence of peritubary adhesions and the nature of their distribution.
- Ultrasonic examination of small pelvis organs - diagnostics of changes in uterine tubes.

In the follow-up postoperative period, colpocytology, hysterosalpingography and hysterosalpingoscopy are recommended for the diagnosis of reproductive disorders, allowing to differentiate the tube-peritoneal form of infertility from the other forms.

Conclusion: thus, prevalent appendicular peritonitis is a severe surgical disease of the abdominal cavity. Patients who have had undergone prevalent appendicular peritonitis in their childhood should be followed-up by pediatric surgeon and a pediatric gynecologist. He proposed algorithm of diagnosis and treatment of PAP complications makes it possible to carry out timely diagnostics and treatment of the revealed reproductive system pathology. In spite of the fact that information concerning the methods of examination of gynecological patients have been presented in a number of manuals [10,11,12], we consider it to be reasonable to apply the proposed algorithm in the management tactics in patients, who underwent prevalent appendicular peritonitis, by supplementing new information and practical recommendations.

References

- [1] Agzamova M.N. The ways of improvement the results of treatment of peritonitis, taking into account the phase of the disease, the degree of contamination of the microflora of the abdominal cavity and immunity: dis. abstract. ... cand. medical. sciences.-Tashkent. 2001.-p.22.
- [2] Bashar X.Sh. Preventive therapy and intraoperative prophylaxis of the complicated course of acute appendicitis in children (clinical and experimental study): Dis. abstract. ... cand. medical sciences.-M.2002.-p.19.
- [3] Yusupov Sh. A., Shamsiyev A.M., Gynecological status of patients who suffered appendicular peritonitis in childhood // Scientific and medical reviewed journal "Avicenna Bulletin" 2020

[4] Polyakova O.V. Surgical treatment of tubal-peritoneal infertility, prevention of the formation of adhesions. Dis. abstract. ...cand. medical sciences. -Ryazan. 2004.-p.18

[5] Sleptsov A.A., Savvina V.A., Varfolomeev A.R., Nikolaev V.N., Petukhov E.I., Zuev A.L., Erdineev T.E. To the treatment of appendicular peritonitis in children // Pediatric Surgery. 2017. Vol. 21. No. 6. pp. 316-320.

[6] Surgical diseases of childhood / Ed. Yu. F. Isakova. - M.: Publishing house "Geotar-Med", 2004. – Vol. 1. – p.630 (233).

[7] Shamsiyev A.M., Yusupov Sh. A. Evaluation of the effectiveness of treatment of common appendicular peritonitis in children/ Journal "Bulletin of the doctor" 2019 №3 Samarkand

[8] Shmyreva E.S., Shapkin V.V., Shapkina A.N. Complex treatment of children with appendicular peritonitis using laparotomic ultrasound // Pacific Medical Journal. 2012. No. 3. pp. 80-82.

[9] Alloo J., Gerstle T. Appendicitis in children less than 3 years of age: a 28-year review //Pediatr Surg Int. - 2004. - Vol.19, №12. – P. 777-779. (276)

[10] Beck D. E., Cohen Z. A prospective, randomized, multicenter, con-trolled study of the safety of Seprafilm adhesion barrier in abdominopelvic surgery of the intestine // Dis Cokon Rectum. – 2003 – Vol.46, №10. - P. 1309-1310. (285)

[11] Müller G., Boman J., Shrier I., Gordon P. Natural history of patients with adhesive small bowel obstruction // Br J Surg. – 2000. – Vol.87, №9 P. 1240-1247.

[12] Sookram J, Naidoo N, Cheddie S. Perforated caecal duplication cyst presenting as an appendicular abscess //S Afr J Surg. 2016 Sep;54(3):42.

[13] Шамсиев М.А, Атакулов Д.О, Юсупов Ш.А. Экспериментальное изучение влияния озона на течение перитонита и спайкообразование // Журнал “Детская хирургия”- 2000. - Т.6 - С. 22-25

[14] Shamsiev A.M, Yusupov Sh.A, Shahriev A.K. Ефективність ультразвукової сонографії при апендикулярних перитонітах у дітей // Журнал “Здобутки клінічної і експериментальної медицини” – 2016. №2. –С. 38-43.

[15] Yusupov Sh.A, Kurbaniyazov Z.B, Davlatov S.S, Rakhmanov K.E, Daminov F.A. Отдаленные результаты оперативного лечения узловых образований щитовидной железы // Журнал “Здобутки клінічної і експериментальної медицини” -2017. №1. – С. 29-35.

[16] Юсупов Ш.А, Исламова Д.С, Гаффаров У.Б. Структура патологии гастродуоденальной зоны у детей с дискинезией желчного пузыря // Журнал “Ученые записки Орловского государственного университета” – 2014. –Т.2, №7. –С. 73-74.

[17] Бердиярова Ш.Ш, Юсупов Ш.А, Назарова Г.Ш. Клинико-лабораторные особенности хронического гематогенного остеомиелита //

Журнал "Central Asian Research Journal for Interdisciplinary Studies (CARJIS)" – 2022. – Т. 2, №5. – С. 116-125.

[18] Юсупов Ш.А, Шамсиев А.М, Атакулов Ж.О, Джалолов Д.А. Оценка интенсивности синдрома эндогенной интоксикации у детей с распространенным аппендикулярным перитонитом // Журнал "Медицинский альманах" – 2019. № 5-6(61) . – С. 57-61.

[19] Шамсиев А.М, Юсупов Ш.А, Шарипов Р.Х. Влияние озонотерапии на показатели перекисного окисления липидов у детей с распространенными формами аппендикулярного перитонита // Журнал "Анналы хирургии" – 2001. – Т. 5. – С. 77.

[20] Юсупов Ш.А, Шамсиев Ж.А, Суванкулов У.Т, Данияров Э.С. Хирургическая тактика при обструктивном калькулезном пиелонефрите у детей // Журнал "Саратовский научно-медицинский журнал" – 2007. – Т. 3, №2. – С. 79-80.

[21] Юсупов Ш.А, Мардыева Г.М, Бахритдинов Б.Р. Особенности рентгенологической семиотики при пневмонии у детей раннего возраста // Журнал "Актуальні питання педіатрії, акушерства та гінекології" – 2017. №2. – С. 21-24.

[22] Шамсиев А.М, Юсупов Ш.А, Юлдашев Б.А, Мухамадиева Л.А. Состояние иммунного статуса у детей с хроническим бронхитом // Журнал "Педиатрический вестник Южного Урала" – 2017. №1. – С. 84-89.

[23] Шамсиев А.М, Юсупов Ш.А, Махмудов З.М. Хирургическое лечение детей с острым гематогенным остеомиелитом костей, образующих тазобедренный сустав // Журнал "Российский вестник детской хирургии, анестезиологии и реаниматологии" – 2014. – Т. 4, №3. – С.86-89.

[24] Юсупов Ш.А, Курбаниязов З.Б, Зайниев А.Ф. Узловые образования щитовидной железы. состояние проблемы (обзор литературы) // Источник "Вісник наукових досліджень" – 2018. №1. – С. 10-15.

[25] Шамсиев А.М, Саидов М.С, Аипов Р.Р, Атакулов Д.О, Юсупов Ш.А. Хирургическая коррекция недержания кала при свищах в половую систему у девочек // Журнал "Российский вестник детской хирургии, анестезиологии и реаниматологии" – 2014. – Т. 4, №2. – С.25-29.

[26] Шамсиев А.М, Саидов М.С, Атакулов Д.О, Юсупов Ш.А, Шамсиев Ж.А, Суванкулов У.Т. Хирургическое лечение аноректальных пороков у детей // Журнал "Вестник хирургии имени ИИ Грекова" – 2011. – Т. 170, №2. – С. 40-43.

[27] Akhmedov I, Khamidov A, Kholmirezayev S, Yusupov Sh, Umarov I. Improving river sediment distribution calculation in mountain rivers // "Science and innovation" V.1 № A8. - P. 1014-1019

[28] Юсупов Ш.А. Диагностическая значимость ультразвуковой сонографии при аппендикулярных перитонитах у детей // Журнал "Сибирский медицинский журнал" (Иркутск) – 2009. – Т. 86, №3. – С. 138-141.

[29] Махмудова, А.Н. and Махмудова, С., 2022. Гуманитаризация медицинского образования как фактор повышения качества обучения в вузе. *Science and Education*, 3(6), pp.709-718.

[30] Махмудова, А.Н., 2022. Правовая защита пациентов в сфере здравоохранения в новом Узбекистане. *Academic research in educational sciences, (Conference)*, pp.102-107.

[31] Махмудова АН, Камариддинзода АК. Защита прав пациентов в Республике Узбекистане. *Science and Education*. 2022;3(10):54-62.

[32] Nugmanovna MA, Kamariddinovna KA. Modern biotechnical problems of medicine and their solutions. In *Archive of Conferences 2021 Jan 28 (Vol. 13, No. 1, pp. 169-173)*.

[33] Kamariddinovna KA, Nugmanovna MA. Improving population health the important task of the state. In *Archive of Conferences 2021 Mar 30 (Vol. 17, No. 1, pp. 204-208)*.

[34] Nugmanovna MA. BIOETHICS AS A FORM OF PROTECTION OF INDIVIDUALITY AND PERSONALIZED MEDICINE. *Thematics Journal of Social Sciences*. 2022 Oct 28;8(4).

[35] Nugmanovna MA. Bioetika zamonaviy madaniyatda individuallikni himoya qilish shakli sifatida. *философия и жизнь международный журнал*. 2022 Nov 30(SI-2).

[36] Махмудова, А.Н., Ибрагимова, Э.Ф., Шукурова, Д.Б., Абдурахмонова, З.Э. and Наимова, З.С., 2020. Медицина Узбекистана-достижения и перспективы развития сферы. *Достижения науки и образования*, (3 (57)), pp.49-52.